Types	Biofilm component	Biofilm phase	State of Development&	Pros	Cons
Agents					
Antibiotics <sup>1,2</sup>	Microbial cell	All stages	Clinical	*Well understood.  *Novel combinations promising.  *Many can be combined with local delivery.	*Resistance.  *Cytotoxicity.  *Not necessarily effective against dormant populations.  *Some have limited penetration into biofilm EPS
Antimicrobial peptides <sup>3</sup>	Microbial cell	All stages	Pre-clinical	*Small molecules easily engineered for optimization.  *Membrane physical disruption reduces probability of resistance.  *Broad-spectrum  *Species-specific targeting possible.	*Charge may limit transport through biofilm EPS. *Potential proteolytic degradation. *pH may affect activity *Delivery to infected site
Antimicrobial oligonucleotides <sup>4,5</sup>	Microbial cell	Early/Mature biofilm	In vivo	*Small molecules easily engineered for optimization.	*Charge may limit transport through biofilm EPS. *Potential binding with eDNA. *Delivery to infected site *Potential degradation by nucleases
Nanoparticles (inorganic, organic, hybrid) <sup>6-8</sup>	Microbial cell, EPS	All stages	In vivo, pre- clinical, clinical†	*Readily functionalized. *Intrinsic bioactivity combined with drug- delivery capacity. *Small size allows transport into the EPS. *Triggered (pH, O <sub>2</sub> ) mechanism possible for	*Charge may limit penetration into the biofilm EPS.  *Properties affected by biological fluids.  *Delivery to the infected site *Cytotoxicity.

on demand treatment.

Other antimicrobials/oxidizers/ antiseptics <sup>1</sup>	Microbial cell	All stages	Clinical	*Physical mode of action not requiring cellular activity. *Broad-spectrum.	*Lack of targeting specificity.  * Restricted transport into biofilm EPS.  *Cytotoxicity.  *Reactive species neutralized by EPS.
Persisters/dormant cells targeting <sup>9</sup>	Microbial cell	Early/Mature biofilm	In vivo	*Specifically targeted to non-growing populations.	*Resistance (not well understood). *Delivery to infected site and transport into the biofilm.
Antibody/Vaccines <sup>1</sup>	Microbial cell, EPS	Initial attachment, early biofilm	In vivo	*Targeted to specific pathogens.	*Restricted transport into biofilm EPS. *Strain replacement. *Disruption of commensal populations.
Adhesin inhibitors/binding <sup>10</sup>	EPS	Initial attachment	Pre-clinical	*Prevention preferable to treatment.	*Symptomatic infections have established biofilms. *Interaction with host components. *Delivery to at risk or infected site.
Bacteriophages <sup>7</sup>	Microbial cell	Early/Mature biofilm	In vivo	*Highly specific and small size to enter biofilm EPS.	*Strain replacement. *Delivery to infected site and transport into the biofilm.
Detergent/Surfactant irrigants <sup>1,11</sup>	Microbial cell, EPS	All stages	Clinical	*Disruption not dependent of killing cells.	*Not all biofilm removed. *Release of pathogens may result in recolonization and

				*Active on dormant cells. *Readily combined for multimodal therapeutics.	acute infection.
Dispersal Inducers <sup>12,13</sup>	Microbial cell	Mature biofilm	In vitro, In vivo, pre-clinical, clinical	*Manipulating natural processes might be less likely to develop resistance.	*Release of pathogens may result in recolonization and acute infection.  *Only portions of the biofilms are released.  *Cytotoxicity.  *Delivery to infected site and transport into the biofilm.
Degradative Enzymes <sup>14,15</sup>	EPS	Early/Mature biofilm	Clinical, pre- clinical	*Disruption not dependent on killing cells.  *Weaken biofilm physical structure; facilitate mechanical removal/mass transport.  *Disrupt pathogenic microenvironment.  *Cell activity not required.  *Readily combined with irrigants and shear.	*Not all biofilm removed, possibly due to complex EPS chemistry and physical structure.  *Release of pathogens may result in recolonization and acute infection.  *Delivery to infected site  *No, or limited, antimicrobial activity.  *Cytotoxicity
EPS synthesis inhibitors <sup>1</sup>	EPS	Initial attachment, early biofilm	In vivo, In vitro	*Prevention of early biofilm formation and EPS protection. *Readily combined with antimicrobials	*Most infections have established biofilms by the time they are symptomatic. *EPS chemistry and structure highly complex. *Delivery to at risk or infected site.

Natural products <sup>16</sup>	Microbial cell, EPS	All stages	In vivo, clinical	*Selected for broad- range of bioactivity (from enzyme inhibitors to antimicrobials). *Chemical diversity with drug-like properties *Multi-mode of action	*Resistance.  *Complex chemistry and isolation procedures.  *Chemical composition variability.  *Target identification  *Cytotoxicity.
Photodynamic substances <sup>17</sup>	Microbial cell	Early/Mature biofilm	In vivo	*Controlled bioactivation options. *On demand activity.	*Light source access required *Delivery of materials to infected site and transport into biofilm. *EPS may protect cells deeper down.
Metabolic interference <sup>12</sup>	Microbial cell	Early/Mature biofilm	In vivo, In vitro	*Community manipulation against pathogens. *Disrupt pathogenic environment (pH). *Manipulating metabolism less likely to develop resistance. *Can trigger disassembly	*Requires specific metabolizing microbes. *Substrate delivery to and transport into biofilms. *Potential substrate utilization by host. *Release of pathogens may result in recolonization and acute infection.
QS inhibitors <sup>18</sup>	Microbial cell	All stages	Pre-clinical, In vivo	*Manipulating natural pathways less likely to develop resistance. *Biofilm inhibition and biofilm dispersal	*Dependent on growth cycle and nutrient source.  *Signals can be washed away or sequestered in the EPS matrix of established biofilm *Complexity of signaling network.

Probiotics <sup>19</sup>	Microbial cell	Initial attachment, early biofilm	In vitro, Pre-clinical (in oral), clinical†	*Community manipulation against pathogens *Concept proven in gut and vaginal biofilms.	* Establishment of probiotic species in oral (and other established) microbiota challenging *Long-term effects unknown
Physical/Electric					
Electric currents/fields <sup>20,21</sup>	Microbial cell, EPS	Early/Mature biofilm	Clinical, pre- clinical	*Projected through induction or connected wires. *On demand antimicrobial generation. *Also promote wound healing.	*Electrochemistry of body fluids not well understood. *Heating of tissue. *Delivery of fields and currents to deep tissue. *Cytotoxicity.
Transducer/pressure waves <sup>22</sup>	Microbial cell, EPS	Early/Mature biofilm	In vivo, pre- clinical	*Readily projected through skin and soft tissue. *Local delivery. *Physical action reduces probability of resistance.	*Limited targeting.  *Influence of pressure waves on viscoelastic biofilms not well understood.  *Local delivery (i.e. shockwave) limited to small and accessible areas.  *Heating cytotoxic effects.
Interfacial tension <sup>23</sup> (microbubbles/droplets)	Microbial cell, EPS	Early/Mature biofilm	Pre-clinical	*Physical action reduces probability of resistance. *Readily combined with irrigants and shear.	*Accessibility.  *Biofilm viscoelasticity can resist removal.  *Residual cells may remain.
Shear stress <sup>22</sup>	Microbial cell, EPS	Early/Mature biofilm	Clinical	*Physical action reduces probability of resistance. *Readily combined with antimicrobials or nanoparticles.	*Accessibility. *Biofilm viscoelasticity can resist removal. *Possible spread of biofilm if not used in combination with antimicrobial agents.
Non-thermal (cold) plasma <sup>24</sup>	Microbial cell	Early/Mature biofilm	In vivo	*Antimicrobials generated locally. *High level of	*Accessibility of plasma. *Biofilm EPS may protect cells deeper down.

Photothermal activation <sup>24</sup> Delivery Systems	Microbial cell	Early/Mature biofilm	In vitro	oxidation/reactive species renders resistance unlikely.  *Antimicrobial activity can be controlled locally.  *Can be readily combined with surface modifications.	*Response to plasma is species-dependent. *Highly localized.  *Delivery to infected site and transport into biofilm. *Accessibility of light. *Biofilm EPS may protect cells.
Bone cements <sup>25</sup>	Microbial cell	Initial attachment.	Clinical	*High concentrations of	*Antimicrobial cytotoxicity.
Done cements	Microbiai cen	mature biofilm	Cillical	antibiotics maintained at site of local infection for extended periods.  *Prophylactic use.	*Development of resistance.
Rinsing fluid/Irrigators <sup>26,27</sup>	Microbial cell, EPS	Mature biofilm	Clinical	*Can be readily combined with antimicrobial agents.	*Accessibility. *Biofilm viscoelasticity can resist removal
Surfaces <sup>28,29</sup>	Microbial cell	Initial attachment	Clinical	*Prevention more effective than treatment. *Access not required after implantation. *Can be targeted to those surfaces prone of biofilm infection.	*Bacteria have non-specific attachment mechanisms.  *Surfaces masked by dead biofilm or host components.  *Stability of surface coatings.  *Finite antimicrobial reservoir/long-term effects
Nanocarriers (nanoparticles/liposomes) <sup>6</sup>	Microbial cell, EPS	Early/Mature biofilm	In vivo, pre-clinical, clinical	*Readily functionalized. *Small size allows transport into the EPS *Carry/release different drug combinations *Triggered (pH, O <sub>2</sub> ) mechanism possible for on demand drug-release.	*Charge may limit penetration into the biofilm EPS.  *Delivery to the infected site *Properties affected by biological fluids.  *Cytotoxicity *Prolonged retention needed for optimal drug release

&Specifically against biofilms

†used clinically to treat other conditions

δ Clinical - already a licensed product available to patients;

Pre-clinical - currently in human trials

In vivo - currently in animal model

In vitro - encompassing basic (polystyrene plate) to advanced biofilm research (i.e. co-culture, explant tissue, patient samples)

## References

- Lebeaux, D., Ghigo, J.-M. & Beloin, C. Biofilm-related infections: bridging the gap between clinical management and fundamental aspects of recalcitrance toward antibiotics. *Microbiology and Molecular Biology Reviews* **78**, 510-543 (2014).
- Bayramov, D. F. & Neff, J. A. Beyond conventional antibiotics New directions for combination products to combat biofilm. *Adv Drug Deliv Rev*, doi:10.1016/j.addr.2016.07.010 (2016).
- Pletzer, D., Coleman, S. R. & Hancock, R. E. Anti-biofilm peptides as a new weapon in antimicrobial warfare. *Curr Opin Microbiol* **33**, 35-40, doi:10.1016/j.mib.2016.05.016 (2016).
- 4 Michaud, G. *et al.* Overcoming antibiotic resistance in Pseudomonas aeruginosa biofilms using glycopeptide dendrimers. *Chemical Science* **7**, 166-182, doi:10.1039/C5SC03635F (2016).
- Ning, Y. *et al.* Efficient suppression of biofilm formation by a nucleic acid aptamer. *Pathog Dis* **73**, ftv034, doi:10.1093/femspd/ftv034 (2015).
- Zazo, H., Colino, C. I. & Lanao, J. M. Current applications of nanoparticles in infectious diseases. *J Control Release* **224**, 86-102, doi:10.1016/j.jconrel.2016.01.008 (2016).
- Brooks, B. D. & Brooks, A. E. Therapeutic strategies to combat antibiotic resistance. *Advanced drug delivery reviews* **78**, 14-27 (2014).
- 8 Allaker, R. P. & Memarzadeh, K. Nanoparticles and the control of oral infections. *Int J Antimicrob Agents* **43**:95–104 (2014).
- 9 Wood, T. K. Combatting bacterial persister cells. *Biotechnol Bioeng* **113**, 476-483, doi:10.1002/bit.25721 (2016).
- 10 Cozens, D. & Read, R. C. Anti-adhesion methods as novel therapeutics for bacterial infections. *Expert Rev Anti Infect Ther* **10**, 1457-1468, doi:10.1586/eri.12.145 (2012).
- Otzen, D. E. Biosurfactants and surfactants interacting with membranes and proteins: Same but different? *Biochim Biophys Acta* **1859**, 639-649, doi:10.1016/j.bbamem.2016.09.024 (2017).
- McDougald, D., Rice, S. A., Barraud, N., Steinberg, P. D. & Kjelleberg, S. Should we stay or should we go: mechanisms and ecological consequences for biofilm dispersal. *Nat Rev Microbiol* **10**, 39-50, doi:10.1038/nrmicro2695 (2011).
- Barraud, N., Kelso, M. J., Rice, S. A. & Kjelleberg, S. Nitric oxide: a key mediator of biofilm dispersal with applications in infectious diseases. *Curr Pharm Des* **21**, 31-42 (2015).
- 14 Kaplan, J. B. Biofilm matrix-degrading enzymes. *Methods Mol Biol* **1147**, 203-213, doi:10.1007/978-1-4939-0467-9\_14 (2014).

- Pleszczynska, M., Wiater, A., Janczarek, M. & Szczodrak, J. (1-->3)-alpha-D-Glucan hydrolases in dental biofilm prevention and control: A review. *International journal of biological macromolecules* **79**, 761-778, doi:10.1016/j.ijbiomac.2015.05.052 (2015).
- Farha, M. A. & Brown, E. D. Strategies for target identification of antimicrobial natural products. *Nat Prod Rep* **33**, 668-680, doi:10.1039/c5np00127g (2016).
- Yin, R. *et al.* Light based anti-infectives: ultraviolet C irradiation, photodynamic therapy, blue light, and beyond. *Current opinion in pharmacology* **13**, 731-762, doi:10.1016/j.coph.2013.08.009 (2013).
- Brackman, G. & Coenye, T. Quorum sensing inhibitors as anti-biofilm agents. *Curr Pharm Des* **21**, 5-11 (2015).
- 19 Vuotto, C., Longo, F. & Donelli, G. Probiotics to counteract biofilm-associated infections: promising and conflicting data. *Int J Oral Sci* **6**, 189-194, doi:10.1038/ijos.2014.52 (2014).
- Nodzo, S. R. *et al.* Cathodic Voltage-controlled Electrical Stimulation Plus Prolonged Vancomycin Reduce Bacterial Burden of a Titanium Implant-associated Infection in a Rodent Model. *Clin Orthop Relat Res* **474**, 1668-1675, doi:10.1007/s11999-016-4705-7 (2016).
- Banerjee, J. *et al.* Silver-zinc redox-coupled electroceutical wound dressing disrupts bacterial biofilm. *PLoS One* **10**, e0119531, doi:10.1371/journal.pone.0119531 (2015).
- Peterson, B. W. *et al.* Viscoelasticity of biofilms and their recalcitrance to mechanical and chemical challenges. *FEMS Microbiol Rev* **39**, 234-245, doi:10.1093/femsre/fuu008 (2015).
- Gomez-Suarez, C., Busscher, H. J. & van der Mei, H. C. Analysis of bacterial detachment from substratum surfaces by the passage of airliquid interfaces. *Applied and environmental microbiology* **67**, 2531-2537, doi:10.1128/AEM.67.6.2531-2537.2001 (2001).
- Miquel, S., Lagrafeuille, R., Souweine, B. & Forestier, C. Anti-biofilm Activity as a Health Issue. *Front Microbiol* **7**, 592, doi:10.3389/fmicb.2016.00592 (2016).
- Howlin, R. P. *et al.* Antibiotic-loaded synthetic calcium sulfate beads for prevention of bacterial colonization and biofilm formation in periprosthetic infections. *Antimicrob Agents Chemother* **59**, 111-120, doi:10.1128/AAC.03676-14 (2015).
- 26 Rmaile, A. *et al.* An experimental and computational study of the hydrodynamics of high-velocity water microdrops for interproximal tooth cleaning. *J Mech Behav Biomed Mater* **46**, 148-157, doi:10.1016/j.jmbbm.2015.02.010 (2015).
- Urish, K. L., DeMuth, P. W., Craft, D. W., Haider, H. & Davis, C. M., 3rd. Pulse lavage is inadequate at removal of biofilm from the surface of total knee arthroplasty materials. *The Journal of arthroplasty* **29**, 1128-1132, doi:10.1016/j.arth.2013.12.012 (2014).
- Besinis, A., De Peralta, T., Tredwin, C. J. & Handy, R. D. Review of nanomaterials in dentistry: interactions with the oral microenvironment, clinical applications, hazards, and benefits. *ACS Nano* **9**, 2255-2289, doi:10.1021/nn505015e (2015).
- Hall-Stoodley, L., Costerton, J. W. & Stoodley, P. Bacterial biofilms: from the natural environment to infectious diseases. *Nat Rev Microbiol* **2**, 95-108, doi:10.1038/nrmicro821 (2004).